

Retinal Detachment Due To Gyrate Atrophy

Gyrate Atrofiye Bağlı Gelişen Retina Dekolmanı

Orhan Ayar¹, Nesibe Akyürek², Mehmet Orçun Akdemir¹, Serpil Yazgan¹

¹ Bülent Ecevit University, Faculty of Medicine Department of Ophthalmology, Zonguldak, Turkey

² Necmettin Erbakan University, Faculty of Medicine, Division of Pediatric Endocrinology and Diabetes, Konya, Turkey

ABSTRACT

In this case study we report a 7-year old child who was first diagnosed with total retinal detachment and later with Gyrate Atrophy. The patient applied to our clinic with visual loss on his right eye without any history of trauma. The biomicroscopic evaluation funduscopy showed a total retinal detachment on the right eye. A retinal detachment surgery was performed on the right eye. After the surgery re-detachment due to a retinal hole occurred, and the patient lost the light sense. The patient was considered inoperable.

Key Words: Gyrate Atrophy, Retinal Detachment

ÖZET

Bu olguda daha önce Gyrate atrofi tanısı almış 7 yaşında bir çocuk hastada gelişen total retina dekolmanı sunulmuştur. Herhangi bir travma öyküsü olmayan hasta sağ gözde görme kaybı ile kliniğe başvurdu. Yapılan muayenesinde sağ gözde retina dekolmanı tespit edildi. Sağ göze retina dekolman cerrahisi uygulandı. Sonrasında retinal delik nedeni ile tekrar dekolman olan hastada ışık görme hissi kaybolduğu için tekrar cerrahi düşünülmeydi.

Anahtar Sözcükler: Gyrate Atrophy, retina dekolmanı

INTRODUCTION

Gyrate atrophy (GA) is a rare, progressive metabolic choroidal and retinal degeneration resulting from the inherited deficiency of the pyridoxal phosphate-dependent mitochondrial matrix enzyme ornithine- δ -aminotransferase (OAT) mapped on chromosome 10q26. OAT deficiency causes hyperornithinemia, which results in progressive chorioretinal atrophy (1).

CASE REPORT

A 7 year-old child presented with decreased vision in his right eye. There was no history of trauma. The patient was diagnosed with GA with decreased night vision, high myopia and characteristic chorioretinal lesions. At presentation he was just perceiving light on right eye, and the best corrected visual acuity was 20/30 with a highly myopic correction (-12,75 D) on left eye. Biomicroscopic examination revealed an absence of red reflex on the right eye.

Address for Correspondence / Yazışma Adresi: Orhan Ayar, MD Bülent Ecevit University, Faculty of Medicine Department of Ophthalmology, 67600 Kozlu Zonguldak, Turkey Tel: 0 372 261 20 00/2265 Fax: 0 372 261 02 64 E-mail: orhanayar@gmail.com

©Telif Hakkı 2015 Gazi Üniversitesi Tıp Fakültesi - Makale metnine <http://medicaljournal.gazi.edu.tr/> web adresinden ulaşılabilir.

©Copyright 2015 by Gazi University Medical Faculty - Available on-line at web site <http://medicaljournal.gazi.edu.tr/>

doi:<http://dx.doi.org/10.12996/gmj.2015.23>

Fundoscopy showed total retinal detachment on the right eye and severe chorioretinal atrophy with characteristic punched-out lesions in the peripheral retina confirming the diagnosis of GA (Figure 1). The plasma ornithine level was 580,73 $\mu\text{mol/L}$ (10-163). The patient had no family history. Genetic testing was not performed in this case but we think the clinical appearance was sufficient for the diagnosis. The patient has undergone vitreoretinal surgery including pars plana vitrectomy, endolaser photocoagulation and intravitreal silicon tamponade. On the follow up examination, re-detachment due to a retinal hole occurred, and the patient lost the light sense. Cataract formation has occurred after the surgery (Figure 2). Afterwards, the patient was considered inoperable.




Figure 1. Fundus photo of the left eye showing chorioretinal atrophy with characteristic lesions in the peripheral retina.




Figure 2. Cataract formation in the right eye after pars plana vitrectomy.

DISCUSSION

This case was reported because the retinal detachment due to Gyrate Atrophy is a rare condition among young children. GA is characterized by hyperornithinemia, typical retinal and choroidal lesions, and the early cataract formation. Patients with GA generally present with decreased night vision and high myopia during adolescence. Early diagnosis in these patients allows for early treatment (1,2). OAT gene studies in children with a high degree of myopia, especially in cases in which there is a complaint of nyctalopia, would be helpful for early identification (3). Retinal detachment occurs infrequently, with only isolated cases in older patients (4,5). In the treatment of GA, a low-protein diet is recommended. An arginine-free diet, may decrease ornithine to normal levels. Lysine supplement may improve the effect of the arginine-free diet (6). However, the long-term effects of this treatment approach have not been completely evaluated.

Conflict of Interest

No conflict of interest was declared by the authors.

REFERENCES

1. Takki KK, Milton RC. The natural history of gyrate atrophy of the choroid and retina. *Ophthalmology* 1981; 88:292-301.
2. Simell O, Takki K. Raised plasma-ornithine and gyrate atrophy of the choroid and retina. *Lancet* 1973;1:1031-3.
3. Kim SJ, Lim DH, Kim JH, Kang SW. Gyrate atrophy of the choroid and retina diagnosed by ornithine- δ -aminotransferase gene analysis: a case report. *Korean J Ophthalmol* 2013;27:388-91.
4. Engelmann K, Richard G, Kohlschütter A, Bialasiewicz AA. [Rhegmatogenous retinal detachment in atrophia gyrate: pars plana vitrectomy with silicone oil instillation]. *Klin Monbl Augenheilkd* 1996;209:373-5.
5. Barnett CP, Lam WC, Schulze A. Retinal detachment causing unilateral blindness in a 12-year-old girl with gyrate atrophy. *J Inherit Metab Dis* 2009;32:670-1.
6. Valle D, Walser M, Brusilow S, Kaiser-Kupfer MI, Takki K. Gyrate atrophy of the choroid and retina: biochemical considerations and experience with an arginine restricted diet. *Ophthalmology* 1981; 88:325-30.