



Estimation of the capacity of emergency surgery in Konya: Nine-year multicenter study

Konya'da acil cerrahi kapasitesinin tahmini: Dokuz yıllık çok merkezli çalışma

Tevfik Küçükkartallar¹, Murat Çakır¹, Ahmet Tekin¹, Mehmet Balasar², Adil Kartal¹, Hande Köksal³, Bülent Erengül⁴, Emin Türk⁵

ABSTRACT

Objective: Although the number of surgical emergencies continues to increase, comprehensive data on emergency surgical admissions are scarce. The aim of this multicenter study was to evaluate the causes, management, and outcomes of the general surgical emergencies in the city of Konya, Turkey.

Material and Methods: The relevant details of the cases admitted and considered to be general surgical emergencies in Konya over a nine-year period (January 2003–January 2012) were analyzed. All demographic data were analyzed statistically.

Results: The study group comprised 21954 cases from 4 hospitals in Konya: 7154 from Konya Numune Hospital, 6,654 from Konya Education and Research Hospital, 6,400 from Necmettin Erbakan University Meram Medical Faculty, and 1,390 from Başkent University Konya Education and Research Hospital. Their mean age was 59.6 years, and the average hospitalization time was 3.3 days. The diagnoses of the admitted patients were as follows: acute appendicitis (59.57%), bowel obstruction (11.12%), trauma (7.97%), strangulated inguinal hernia (5.46%), acute cholecystitis (4.87%), peptic ulcer perforation (4.09%), mesenteric ischemia (2.73%), necrotizing fasciitis (2.73%), gastrointestinal system bleeding (1.79%), and others (1.1%).

Conclusion: The findings of the study indicate a steady increase in surgical admissions to emergency units. Non-traumatic acute abdomen was the most common reason for general surgical emergencies. Although the number of elderly patients increased, the hospital stay and mortality rates decreased over the study period.

Keywords: Emergency, hospitalization, surgery

ÖZ

Amaç: Cerrahi acillerin sayısının sürekli artmasına rağmen acil cerrahi yatışları hakkında kısıtlı veri bulunmaktadır. Bu çok merkezli çalışmanın amacı, Konya şehrinde acil genel cerrahi sonuçlarını değerlendirmektir.

Gereç ve Yöntemler: Konya'da tüm genel cerrahi acil başvurularının dokuz yıllık sonuçları incelendi (Ocak 2003-Ocak 2012). Tüm demografik veriler istatistiksel olarak değerlendirildi.

Bulgular: Konya'da yer alan 4 hastaneden 21954 hasta çalışmaya alınmıştır. Konya Numune Hastanesi (7154), Konya Eğitim ve Araştırma Hastanesi (6654), Necmettin Erbakan Üniversitesi Meram Tıp Fakültesi (6400) ve Başkent Üniversitesi Konya Eğitim ve Araştırma Hastanesi (1390) verileri toplandı. Hastaların yaş ortalaması 59,6 ve ortalama hastanede kalış süresi 3,3 gündü. Hastaneye kabul edilen hastaların tanıları; akut apandisit (%59,57), bağırsak tıkanıklığı (%11,12), travma (%7,97), boğulmuş kasık fıtığı (%5,46), akut kolesistit (%4,87), peptik ülser perforasyonu (%4,09), mezenterik iskemi (%2,73), nekrotizan fasiit (%2,73), gastrointestinal sistem kanaması (%1,79) ve diğerleriydi (%1,1).

Sonuç: Bu çalışma ile cerrahi kabullerin sürekli arttığı görülmüştür. Non-travmatik akut karın, genel cerrahi acillerinin en sık nedeniydi. Yaşlı hastaların sayısında artış olmasına rağmen, hastanede kalış süresi ve mortalite oranları azalmıştır.

Anahtar Kelimeler: Acil, hastaneye yatış, cerrahi

INTRODUCTION

Emergencies requiring surgical intervention are increasing steadily worldwide; yet, comprehensive data on emergency surgical admissions are scarce. Despite the fact that such emergencies have similar patterns throughout the world, they may differ from one hospital to another (1). A statistical analysis of the clinical workload has gained impetus for reasons such as widespread medical audits, the need to assess the causes, management, and outcomes of general surgical emergencies, and the need to understand patient demographic and population-based profiles. Although surgical emergencies are an essential part of a hospital's workload, reports regarding the increase in emergency unit admissions (2) are either related to overall or medical admissions (3, 4).

The study by Chezian et al. (5) also reported an increase in emergency admissions between 1992 and 1999; however, the cases referred to surgery were beyond the scope of their study. Likewise, to the best of our knowledge, such cases have not been the subject of any report or have been documented after a long-term study. Increases in the aging population, social deprivation, and awareness are associated with the increase in emergency admission rates (2-4).

The aim of the present multicenter study was to assess the general surgical emergency causes, their management, and outcomes in the city of Konya, Turkey.

¹Department of General Surgery, Necmettin Erbakan University Meram School of Medicine, Konya, Turkey

²Department of Urology, Necmettin Erbakan University Meram School of Medicine, Konya, Turkey

³Clinic of General Surgery, Konya Numune Hospital, Konya, Turkey

⁴Clinic of General Surgery, Konya Training and Research Hospital, Konya, Turkey

⁵Department of General Surgery, Başkent University Konya Training and Research Hospital, Konya, Turkey

Address for Correspondence Yazışma Adresi

Murat Çakır

e-mail: drmuratcakir@hotmail.com

Received / Geliş Tarihi: 23.02.2015

Accepted / Kabul Tarihi: 20.06.2015

Available Online Date /

Çevrimiçi Yayın Tarihi: 27.10.2016

©Copyright 2016

by Turkish Surgical Association

Available online at

www.ulusalcerahidergisi.org

©Telif Hakkı 2016

Türk Cerrahi Derneği

Makale metnine

www.ulusalcerahidergisi.org

web sayfasından ulaşılabilir.

MATERIAL AND METHODS

Surgical emergency admissions covering a nine-year period (January 2003-January 2012) were analyzed retrospectively. Hereby, the data obtained from the four participating hospitals- Konya Numune Hospital, Konya Training and Research Hospital, Necmettin Erbakan University Meram School of Medicine, and Bařkent University Konya Training and Research Hospital-were analyzed.

All patients included in this study presented to the emergency service and were hospitalized in a general surgery clinic in Konya. Information on the work carried out was obtained from records of the monthly meetings held in Konya that have been regularly carried out for 15 years to discuss emergency cases. Data regarding patient evaluation, number of hospitalized patients, as well as age, diagnosis, and length of hospital stay were included in the study. The information was collected in accordance with the Declaration of Helsinki.

Statistical Analysis

Statistical analyses were performed using the Statistical Package for the Social Sciences 22.0 (IBM Corp.; Armonk, NY, USA) software package.

RESULTS

In total, 21,954 patients were included from the 4 participating hospitals in Konya: 7,154 from Konya Numune Hospital, 6,654 from Konya Training and Research Hospital, 6,400 from Necmettin Erbakan University Meram School of Medicine, and 1,390 from Bařkent University Konya Training and Research Hospital (Figure 1).

Admission rates were higher among the elderly at Necmettin Erbakan University Meram School of Medicine as compared to other centers. The majority of the patients admitted were male, and were residents of Konya (77.7%). The participants' mean age was 59.6 years, with a mean hospital length of stay of 3.3 days. The mean delay from the beginning of the symptoms until presentation to the emergency service was 2.1 days. The diagnoses of the admitted patients were as follows: acute appendicitis (59.57%), bowel obstruction (11.12%), trauma (7.97%), strangulated inguinal hernia (5.46%), acute cholecystitis (4.87%), peptic ulcer perforation (4.09%), mesenteric ischemia (2.73%), necrotizing fasciitis (2.73%), gastrointestinal system (GIS) bleeding (1.79%), and others (1.1%) (Figure 2a-d).

The mean age of the patients admitted to the university hospital emergency clinics was higher than in the other hospitals. Most patients admitted to all four hospitals were diagnosed as having appendicitis, but in the university hospitals, the ratio of appendicitis was lower than that of other hospitals. In contrast, most of the complicated cases (75%) were admitted to the university hospitals.

While the number of open cholecystectomies for acute cholecystitis was higher in the first few years of this study, with the increase in the laparoscopic cholecystectomy rate and surgeon experience, this rate dropped in the latter few years. Likewise, for gastrointestinal bleeding requiring surgical intervention, with advancements in medical treatment, the need for surgery significantly decreased. According to this data, patients with diseases that have high morbidity and mortality rates and re-




Figure 1. Patient demographic data




Figure 2. a-d. (a) Emergency Surgical admissions in Necmettin Erbakan University Meram Medical Faculty. (b) Emergency Surgical admissions in Konya Education and Research Hospital. (c) Emergency Surgical Admissions in Konya Numune Hospital. (d) Emergency Surgical Admissions in Bařkent University Konya Training and Research Hospital

quire longer hospital care and follow-up, such as necrotizing fasciitis and mesenteric ischemia, were only accepted to university hospitals. While the average age of the patients admitted to the university hospitals increased, the average hospital stay decreased as a result of advancements in postoperative care, equipment, and medication.

DISCUSSION

The progressive increase in the number of emergency surgical admissions and decreasing length of hospital stay shown in this study were in line with other recent reports on the overall tendencies in emergency units (2). These might be attributed partially to the increasing population and partially to the rapid increase in the elderly population specifically (6, 7). However, these are not the only reasons for the increase. Currently, people are presenting to emergency units more. This might be the underlying reason for the steady increase in the high number of abdominal pain and constipation cases in the emergency units. Moreover, to some extent, a particular number of emergency admissions are re-admissions (perhaps due to early discharge from the hospital). However, these cases are outside the scope of this study. The increasing number of emergency unit admissions did not result in an apparent need for more surgical beds, contrary to the observations made by Chezian et al. (5). Nevertheless, the increased workload in the emergency wards has brought about increased nursing, medical, and auxiliary staff needs. Bagust et al. (8) highlighted the extra empty-bed capacity needed for effective emergency unit admission management. The roots of the perceived crises within the health care system are attributed to these important considerations.

In a recent study, acute appendicitis was the most common surgical emergency (1, 9). This finding is in concordance with reports from different parts of the world with different patient series, including pregnant women. However, Anyanwu et al. (10) reported in their 1999 study conducted at the University of Ilorin Teaching Hospital, Nigeria, that superficial skin trauma was the most frequent reason for emergency surgery, followed by intestinal obstruction and appendicitis. In the present study, while appendicitis was the most frequent surgical emergency in non-university hospitals, more complicated cases, such as ileus, penetrating injuries, mesenteric ischemia, and necrotizing fasciitis were mostly admitted to the university hospitals.

Furthermore, a patient's age was an important determinant of the frequency and outcome of abdominal surgical emergencies. In the elderly with severe systemic disorders, abdominal emergency surgery can be a life-threatening condition, and hence requires a more careful evaluation as compared to younger patients. Although the number of elderly patients admitted increased between 2003 and 2012, the total number of beds occupied by elderly patients actually decreased due to shorter hospital stay. However, it would be difficult to shorten the average hospital stay any further. In this study, the mean hospital stay was 3.3 days, while it was 5.5 days in the elderly (i.e., over 70 years of age).

In our experience, the mortality rate increase in elderly patients can be attributed to perioperative risks, delay in surgical treatment, conditions that only permit palliative surgery, comorbidities, higher American Society of Anesthesiologists (ASA) grading, age above 80 years, colorectal surgery, malignant diseases, and the severity of the surgical condition.

As reported above, strangulated hernia ranked high among surgical emergencies, with a rate of 10-25% (11-14). In the present study, its frequency was 15% in university hospitals. Most of the patients were aware of the presence of an external hernia long before they became aware of the strangulation. Elective surgical management of abdominal hernia at a convenient time could prevent development of emergencies in most hernia cases. In the present study, the rate of bowel necrosis was low.

Furthermore, advanced age is an important factor contributing to the frequency of intricate malignant neoplasm. Advanced age contributes to the high mortality in free perforations into the peritoneal cavity and abdominal sepsis (6, 15). Mechanical obstructions had a significant part in this study. According to a recent report, the frequency of intestinal obstruction ranged from 15-20% of total surgical emergencies, most of them due to postoperative adhesions. In line with the findings of the present study, obstructive malignancies were reported as an age-related pathology.

In addition, geographical location had a significant effect on large bowel obstruction in this study. Compared to the elderly Western population, volvulus is the leading cause of large bowel obstruction in Turkey. In particular, cardiovascular disorders may lead to mesenteric vascular occlusion that may lead to a surgical emergency (16). The frequency of acute mesenteric ischemia was less than 10% in the present study, similar to other recent reports. Its clinical course and outcomes are more destructive than any other abdominal emergency.

In the present study, vascular occlusion related bowel necrosis ranked the highest among surgical emergencies. The surgical intervention for mesenteric vascular occlusion related bowel necrosis is limited and rarely successful. The findings of Mamode et al. (17) were in line with ours reporting 81% mortality.

Acute gastrointestinal hemorrhage necessitating a surgical intervention has severe consequences, depending on concomitant diseases. Although its frequency is rare, the mortality rate in surgically treated gastrointestinal hemorrhage is the highest as compared to other surgical emergencies. The prognosis of upper gastrointestinal hemorrhage was less evident pronounced in patients with serious comorbidities, with a mortality rate reaching up to 22.4% (6). An increase in operations for biliary and diverticular disease was reported in several studies related to emergency wards (18, 19). Regardless of age, acute calculous cholecystitis is the most common surgery in relation to biliary diseases. In line with the current literature, the present study determined that peptic ulcer complications are now fewer, due to the presence of more effective medical therapies.

CONCLUSION

- The present study documented a steady increase in emergency unit surgical admissions.
- Non-traumatic acute abdomen was the most common reason for general surgery admissions.
- Appendectomy was the most frequent operation in state hospitals.
- Mesenteric ischemia was the most fatal emergency.

Ethics Committee Approval: Authors declared that the research was conducted according to the principles of the World Medical Association Declaration of Helsinki "Ethical Principles for Medical Research Involving Human Subjects", (amended in October 2013).

Informed Consent: Written informed consent was obtained from patients who participated in this study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - A.T.; Design - M.Ç.; Supervision - A.K.; Resources - T.K.; Materials - B.E.; Data Collection and/or Processing - E.T.; Analysis and/or Interpretation - M.B.; Literature Search - M.B.; Writing Manuscript - M.Ç.; Critical Review - M.Ç.

Acknowledgements: We would like to express Dr Adnan Kaynak for his contributions and Dr. Harun Şimşek for his English review.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

Etik Komite Onayı: Yazarlar çalışmanın World Medical Association Declaration of Helsinki "Ethical Principles for Medical Research Involving Human Subjects", (amended in October 2013) prensiplerine uygun olarak yapıldığını beyan etmişlerdir.

Hasta Onamı: Yazılı hasta onamı bu çalışmaya katılan hastalardan alınmıştır.

Hakem Değerlendirmesi: Dış bağımsız.

Yazar Katkıları: Fikir - A.T.; Tasarım - M.Ç.; Denetleme - A.K.; Kaynaklar - T.K.; Malzemeler - B.E.; Veri Toplanması ve/veya İşlemesi - E.T.; Analiz ve/veya Yorum - M.B.; Literatür Taraması - M.B.; Yazıyı Yazan - M.Ç.; Eleştirel İnceleme - M.Ç.

Teşekkür: Uzman katkılarından dolayı Dr. Adnan Kaynak ve İngilizce düzeltmelerden dolayı Dr. Harun Şimşek'e teşekkür etmek istiyoruz.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemişlerdir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

REFERENCES

- Chianakwana GU, Ihegihu CC, Okafor PI, Anyanwu SN, Mbonu OO. Adult surgical emergencies in a developing country: the experience of Nnamdi Azikiwe University Teaching Hospital, Nnewi, Anambra State, Nigeria. *World J Surg* 2005; 29: 804-807. [CrossRef]
- Campbell WB, Lee EJ, Van de, Sijpe K, Gooding J, Cooper MJ. A 25-year study of emergency surgical admissions. *Ann R Coll Surg Engl* 2002; 84: 273-277. [CrossRef]
- Curley PJ, Spark JI, Kester RC, Scott DJA. Audit of vascular surgical workload: use of data for service development. *Ann R Coll Surg Engl* 1996; 78: 209-213.
- Ashraf SQ, Bajwa A, Magee TR, Galland RB. Vascular surgery with-in general surgery: an analysis of workload 1989-2005. *Ann R Coll Surg Engl* 2007; 89: 682-684. [CrossRef]
- Chezian C, Pye J, Jenkinson LR. The next millennium are we becoming emergency surgeons? A seven year audit of surgical and urological admissions in a rural district general hospital. *Ann R Coll Surg Engl* 2001; 83: 117-120.
- Gürleyik G, Gürleyik E, Unalmışer S. Abdominal surgical emergency in the elderly. *Turk J Gastroenterol* 2004; 13: 47-52.
- Breen A, Carr E, Mann E, Crossen-White H. Acute back pain management in primary care: a qualitative pilot study of the feasibility of a nurse-led service in general practice. *J Nurs Manag* 2004; 12: 201-209. [CrossRef]
- Bagust A, Place M, Posnett JW. Dynamics of bed use in accommodating emergency admissions: a stochastic simulation model. *BMJ* 1999; 319: 155-158. [CrossRef]
- Tekin K, Ozek MC. Acute appendicitis after blunt abdominal trauma. *Ulus Travma Derg* 2001; 7: 207-209.
- Anyanwu SN. Abdominal surgical emergencies in a Nigerian teaching hospital. *Orient J Med* 1989; 1: 18-21.
- Rigberg D, Cole M, Hiyama D, McFadden D. Surgery in the nineties. *Am Surg* 2000; 66: 813-816.
- Arhinful E, Jenkins D, Schiller HJ, Cullinane DC, Smoot DL, Zielinski MD. Outcomes of damage control laparotomy with open abdomen management in the octogenarian population. *J Trauma* 2011; 70: 616-621. [CrossRef]
- Agresta F, Mazzarolo G, Ciardo LF, Bedin N. The laparoscopic approach in abdominal emergencies: has the attitude changed? : A single-center review of a 15-year experience. *Surg Endosc* 2008; 22: 1255-1262. [CrossRef]
- Vatansev C, Tekin A. Yaşlılarda Fıtık Cerrahisi Türkiye Klinikleri. *J Surg Med Sci* 2006; 2: 86-88.
- Lewis LM, Banet GA, Blanda M, Hustey FM, Meldon SW, Gerson LW. Etiology and clinical course of abdominal pain in senior patients: a prospective, multicenter study. *J Gerontol A Biol Sci Med Sci* 2005; 60: 1071-1076. [CrossRef]
- Ragsdale L, Southerland L. Acute abdominal pain in the older adult. *Emerg Med Clin North Am* 2011; 29: 429-448. [CrossRef]
- Mamode N, Pickford I, Leiberman P. Failure to improve outcome in acute mesenteric ischaemia: seven-year review. *Eur J Surg* 1999; 165: 203-208. [CrossRef]
- Jaramov N, Sokolov M, Angelov K, Toshev S, Petrov B. Diverticulosis of the colon end its complications under the mask of emergency surgical abdomen--10 years experience. *Khirurgiia (Sofia)* 2009; 1: 5-9.
- Borzellino G, Sauerland S, Minicozzi AM, Verlato G, Di Pietrantonj C, de Manzoni G, et al. Laparoscopic cholecystectomy for severe acute cholecystitis. A meta-analysis of results. *Surg Endosc* 2008; 22: 8-15. [CrossRef]