

# Explicit Solutions of a Three-Dimensional System of Nonlinear Difference Equations

Bahriye Yilmazyıldırım<sup>a</sup>, Durhasan Turgut Tollu<sup>b</sup> 

<sup>a</sup>Necmettin Erbakan University Graduate School of Natural and Applied Science, Turkey

<sup>b</sup>Necmettin Erbakan University, Department of Mathematics-Computer Sciences, Konya, Turkey

## ABSTRACT

In this paper, we show that the system of difference equations

$$x_{n+1} = \frac{x_n + y_n}{1 + x_n y_n}, y_{n+1} = \frac{y_n + z_n}{1 + y_n z_n}, z_{n+1} = \frac{z_n + x_n}{1 + z_n x_n}, n \in \mathbb{N}_0,$$

where the initial values  $x, y, z$  are real numbers, are solvable in explicit form via some changes of variables and tricks. Also, we determine the forbidden set of the initial values  $x, y, z$  for the above mentioned system and investigate asymptotic behavior of the well-defined solutions by using these explicit formulas.

### Keywords:

Asymptotic behavior; explicit solution; rational difference equation; system

### AMS Classification:

39A10

## INTRODUCTION

Nonlinear difference equations constitute an important class of difference equations and studying of this kind of equations have recently attracted great interest. One can see this in some recent studies. See, for example [2-4,7-8,15-16,18,24-27]. Particularly, there have been a renewed interest on solvable ones of such equations and systems. For example, published papers on solvability of some types can be found in the references [5-6,10-14,28,31]. Additionally, there are some equations and systems whose solvability are newly discovered. For example, the solvability of the nonlinear difference equation

$$x_{n+1} = \frac{a + x_n x_{n-1}}{x_n + x_{n-1}}, n \in \mathbb{N}_0, \quad (1)$$

where  $a \in [0, \infty)$  and the initial values  $x_1, x_0$  are real numbers, which was studied by Xianyi and Deming in [29], is newly discovered. The fact remains that the equations and the systems in the references [1,19-21,23,30], are so.

Our aim in this study is to show that the following systems of difference equations

$$x_{n+1} = \frac{x_n + y_n}{1 + x_n y_n}, y_{n+1} = \frac{y_n + z_n}{1 + y_n z_n}, z_{n+1} = \frac{z_n + x_n}{1 + z_n x_n}, n \in \mathbb{N}_0, \quad (2)$$

where real initial values  $x_0, y_0$  and  $z_0$  are real numbers, can be solved in explicit form. Also, we determine the forbidden set of the initial values  $x_0, y_0, z_0$  for the system and investigate asymptotic behavior of the well-defined solutions by using these formulas.

### Definition

Let

$$\begin{aligned} x_{n+1} &= f(x_n, y_n, z_n), y_{n+1} = g(x_n, y_n, z_n), \\ z_{n+1} &= h(x_n, y_n, z_n), n \in \mathbb{N}_0, \end{aligned} \quad (3)$$

where  $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ ,  $g: \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $h: \mathbb{R}^3 \rightarrow \mathbb{R}$  is given functions, be a system of difference equations of first-order, and  $D_f, D_g$  and  $D_h$  be the domains of the functions  $f, g$  and  $h$ , respectively. Forbidden set of difference equation (3) is given by

$$F = \left\{ \begin{aligned} &(x_0, y_0, z_0) \in \mathbb{R}^3 : (x_i, y_i, z_i) \in D \text{ for} \\ &i \in \overline{0, n-1}, n \in \mathbb{N}_0, \text{ and } (x_n, y_n, z_n) \notin D \end{aligned} \right\}$$

### Article History:

Received: 2017/06/21

Accepted: 2017/08/25

Online: 2017/12/22

### Correspondence to:

Durhasan Turgut Tollu

Necmettin Erbakan University, Faculty

of Science, Department of Mathematics-

Computer Sciences, Konya, Turkey

E-Mail: ttollu@konya.edu.tr

where  $D := D_f \times D_g \times D_h$ . This set contains all the initial values which causes the undefinable solutions of the system. That is, the initial values chosen from the complement of the forbidden set always produce the well-defined solutions.

## RESULTS

In this section, we give our main results by obtaining the general solution in explicit form of the system (2). Next, we determine the forbidden set of the initial values  $x_0, y_0, z_0$  for the system (2). Additionally, we investigate asymptotic behavior of the well-defined solutions by using their explicit formulas.

### The Explicit General Solution Of The System

To solve the system (2), we apply the changes of variables

$$x_n = \frac{e^{u_n} - 1}{e^{u_n} + 1}, y_n = \frac{e^{v_n} - 1}{e^{v_n} + 1}, z_n = \frac{e^{w_n} - 1}{e^{w_n} + 1} \quad (4)$$

to the system. Then, we have the linear system

$$u_{n+1} = u_n + v_n, v_{n+1} = v_n + w_n, w_{n+1} = w_n + u_n, n \in \mathbb{N}_0 \quad (5)$$

The system (5) can be written as

$$u_{n+3} - 3u_{n+2} + 3u_{n+1} - 2u_n = 0, n \in \mathbb{N}_0, \quad (6)$$

$$v_{n+3} - 3v_{n+2} + 3v_{n+1} - 2v_n = 0, n \in \mathbb{N}_0, \quad (7)$$

and

$$w_{n+3} - 3w_{n+2} + 3w_{n+1} - 2w_n = 0, n \in \mathbb{N}_0, \quad (8)$$

which are disjoint. Note that the equations (6)-(8) are in the same form. Therefore, we only solve one of them. Let choose Eq. (6) which can be written as

$$(u_{n+3} - 2u_{n+2}) - (u_{n+2} - 2u_{n+1}) + (u_{n+1} - 2u_n) = 0, n \in \mathbb{N}_0.$$

Set  $u_{n+2} - 2u_{n+1} = \tilde{u}_n$ , So, we get the linear equation of order two

$$\tilde{u}_{n+1} = \tilde{u}_n - \tilde{u}_{n-1}, n \in \mathbb{N}_0, \quad (9)$$

Eq. (9) is periodic with period 6 such that  $\tilde{u}_{6n+i} = \tilde{u}_i$ , ( $i = 0, \bar{5}$ ) which implies

$$u_{6n+2+i} - 2u_{6n+1+i} = u_{i+2} - 2u_{i+1}, n \in \mathbb{N}_0. \quad (10)$$

By adding the backward iterate of (10) to own itself, we get

$$u_{6n+2+i} - 2u_{6n+1+i} + u_{6n+1+i} - 2u_{6n+i} = u_{i+2} - 2u_{i+1} + u_{i+1} - 2u_i$$

or

$$u_{6n+2+i} - u_{6n+1+i} - 2u_{6n+i} = u_{i+2} - u_{i+1} - 2u_i, n \in \mathbb{N}_0. \quad (11)$$

Eq. (6) can be also written in the further form

$$(u_{n+3} - u_{n+2} + u_{n+1}) - 2(u_{n+2} - u_{n+1} + u_n) = 0. \quad (12)$$

If we apply the change of variables

$$u_{n+2} - u_{n+1} + u_n = \hat{u}_n \quad (13)$$

to Eq. (12), then, from Eq. (12), it follows that

$$\hat{u}_{n+1} - 2\hat{u}_n = 0 \quad (14)$$

whose the general solution is given by

$$\hat{u}_n = 2^n \hat{u}_0 \quad (15)$$

which implies

$$u_{n+2} - u_{n+1} + u_n = 2^n (u_2 - u_1 + u_0). \quad (16)$$

Eq. (16) can be decomposed in terms of its own subscript as follows;

$$u_{6n+2+i} - u_{6n+1+i} + u_{6n+i} = 2^{6n+i} (u_2 - u_1 + u_0), i = \overline{0,5} \quad (17)$$

By subtracting (11) from (17), we get the formula

$$u_{6n+i} = \frac{1}{3} (2^{6n+i} (v_1 + u_0) - (v_{i+1} - 2u_i)), i = \overline{0,5}, \quad (18)$$

for the solution  $(u_n)$  of Eq. (6). We also state the formula (18) explicitly such that

$$u_{6n} = \frac{2^{6n} + 2}{3} u_0 + \frac{2^{6n} - 1}{3} v_0 + \frac{2^{6n} - 1}{3} w_0, \quad (19)$$

$$u_{6n+1} = \frac{2^{6n+1} + 1}{3} u_0 + \frac{2^{6n+1} + 1}{3} v_0 + \frac{2^{6n+1} - 2}{3} w_0, \quad (20)$$

$$u_{6n+2} = \frac{2^{6n+2} - 1}{3} u_0 + \frac{2^{6n+2} + 2}{3} v_0 + \frac{2^{6n+2} - 1}{3} w_0, \quad (21)$$

$$u_{6n+3} = \frac{2^{6n+3} - 2}{3} u_0 + \frac{2^{6n+3} + 1}{3} v_0 + \frac{2^{6n+3} + 1}{3} w_0, \quad (22)$$

$$u_{6n+4} = \frac{2^{6n+4} - 1}{3} u_0 + \frac{2^{6n+4} - 1}{3} v_0 + \frac{2^{6n+4} + 2}{3} w_0, \quad (23)$$

$$u_{6n+5} = \frac{2^{6n+5} + 1}{3} u_0 + \frac{2^{6n+5} - 2}{3} v_0 + \frac{2^{6n+5} + 1}{3} w_0 \quad (24)$$

for  $n \in \mathbb{N}_0$ , Consequently, by the change of variables  $x_n = \frac{e^{u_n} - 1}{e^{u_n} + 1}$ , we have the formulas

$$x_{6n} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n}+2}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n}-1}{3}} + 1} \quad (25)$$

$$x_{6n+1} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+1}+1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+1}+1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+1}-2}{3}} + 1} \quad (26)$$

$$x_{6n+2} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+2}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+2}-2}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+2}-1}{3}} + 1} \quad (27)$$

$$x_{6n+3} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+3}-2}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+3}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+3}-1}{3}} + 1} \quad (28)$$

$$x_{6n+4} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+4}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+4}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+4}-2}{3}} + 1} \quad (29)$$

$$x_{6n+5} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+5}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+5}-2}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+5}-1}{3}} + 1} \quad (30)$$

which are the formulas of the variable  $x_n$ , for  $n \in \mathbb{N}_0$ , The formulas of  $y_n$  can be obtained by the first equation of (2). That is, by solving  $y_n$  in this equation, we have

$$y_n = \frac{x_{n+1} - x_n}{1 - x_{n+1}x_n} \quad (31)$$

From (25)-(31), it follows that

$$y_{6n} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n}-2}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n}-1}{3}} + 1} \quad (32)$$

$$y_{6n+1} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+1}-2}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+1}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+1}-1}{3}} + 1} \quad (33)$$

$$y_{6n+2} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+2}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+2}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+2}-2}{3}} + 1} \quad (34)$$

$$y_{6n+3} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+3}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+3}-2}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+3}-1}{3}} + 1} \quad (35)$$

$$y_{6n+4} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+4}-2}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+4}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+4}-1}{3}} + 1} \quad (36)$$

$$y_{6n+5} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+5}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+5}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+5}-2}{3}} + 1} \quad (37)$$

for  $n \in \mathbb{N}_0$ , Similarly, from the second equation of (2), we obtain

$$z_n = \frac{y_{n+1} - y_n}{1 - y_{n+1}y_n} \quad (38)$$

which yields

$$z_{6n} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n}-2}{3}} + 1}$$

$$z_{6n+1} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+1}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+1}-2}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+1}-1}{3}} + 1}$$

$$z_{6n+2} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+2}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+2}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+2}-1}{3}} + 1}$$

$$z_{6n+3} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+3}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+3}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+3}-2}{3}} + 1}$$

$$z_{6n+4} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+4}-1}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+4}-2}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+4}-1}{3}} + 1}$$

$$z_{6n+5} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+5}-2}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+5}-1}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+5}-1}{3}} + 1}$$

for  $n \in \mathbb{N}_0$ , along with (32)-(38).

### The Forbidden Set Of The Initial Values

The above obtained formulas exactly determine the solutions of the system (2). But, some initial values yield undefinable solution of the system. Now, we give the set of such initial values by using the formulas. To do this, we use the changes of variables (4) along with the formula (18) and so get the closed formula of  $x_n$  as follows:

$$x_{6n+i} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+x_i}{1-x_i}\right)^2 \left(\frac{1-y_{i+1}}{1+y_{i+1}}\right) + 1} \quad (39)$$

for  $i = \overline{0,5}$  and  $n \in \mathbb{N}_0$ , It is easy to see that the formula (39) is undefinable, if

$$\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+x_i}{1-x_i}\right)^2 \left(\frac{1-y_{i+1}}{1+y_{i+1}}\right) = -1$$

for  $i = \overline{0,5}$  and  $n \in \mathbb{N}_0$ . Similarly, we have the closed formulas of  $y_n$  and  $z_0$  as follows

$$y_{6n+i} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+y_i}{1-y_i}\right)^2 \left(\frac{1-z_{i+1}}{1+z_{i+1}}\right) + 1} \quad (40)$$

and

$$z_{6n+i} = 1 - \frac{2}{\left(\frac{1+x_0}{1-x_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+y_0}{1-y_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+z_0}{1-z_0}\right)^{\frac{2^{6n+i}}{3}} \left(\frac{1+z_i}{1-z_i}\right)^2 \left(\frac{1-x_{i+1}}{1+x_{i+1}}\right) + 1} \quad (41)$$

for  $i = \overline{0,5}$  and  $n \in \mathbb{N}_0$  respectively. Consequently, we find the forbidden set of the initial values  $x_0, y_0, z_0$  as follows:

$$F = \left\{ \begin{array}{l} (x_0, y_0, z_0) \in \mathbb{R}^3 : \\ \left( \frac{1+x_0}{1-x_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+y_0}{1-y_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+z_0}{1-z_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+x_i}{1-x_i} \right)^2 \left( \frac{1-y_{i+1}}{1+y_{i+1}} \right) = -1 \\ \left( \frac{1+x_0}{1-x_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+y_0}{1-y_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+z_0}{1-z_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+y_i}{1-y_i} \right)^2 \left( \frac{1-z_{i+1}}{1+z_{i+1}} \right) = -1, \\ \left( \frac{1+x_0}{1-x_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+y_0}{1-y_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+z_0}{1-z_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+z_i}{1-z_i} \right)^2 \left( \frac{1-x_{i+1}}{1+x_{i+1}} \right) = -1 \\ i = \overline{0, 5}, n \in \mathbb{N}_0 \end{array} \right.$$

**Asymptotic Behavior Of The Well-Defined Solutions**

We investigate the asymptotic behavior of the well-defined solutions of the system (2). The main result of this subsection is given by the following theorem.

**Theorem**

Suppose that the sequence  $(x_n, y_n, z_n)_{n \geq 0}$  is a well-defined solution of the system (2), that is,  $(x_0, y_0, z_0) \notin F$ , Then,

$$\lim_{n \rightarrow \infty} (|x_n|, |y_n|, |z_n|) = (1, 1, 1).$$

**Proof**

To prove, we use the function  $f(x) = \frac{1-x}{1+x}$  along with the formulas (39)-(41). Before proving, we can say that the points  $(0, 0, 0)$ ,  $(1, 1, 1)$  and  $(-1, -1, -1)$  are equilibrium points of the system (2). That is, equilibrium solutions of the system (system) are given by  $(x_n, y_n, z_n)_{n \geq 0} = (0, 0, 0)$ ,  $(x_n, y_n, z_n)_{n \geq 0} = (1, 1, 1)$  and  $(x_n, y_n, z_n)_{n \geq 0} = (-1, -1, -1)$ , respectively. We here deal with nonequilibrium solutions of the system (2). We observe for the function  $f$  that if  $x \in (-\infty, 0)$ , then  $f(x) \in (-1, 1)$ ; if  $x \in (0, 1) \cup (1, \infty)$ , then  $f(x) \in (-\infty, -1) \cup (1, \infty)$ , Hence, we prove the theorem in three cases:

(i) If  $x_0, y_0, z_0 \in (-\infty, 0) \setminus F$ , then

$f(x_0), f(y_0), f(z_0) \in (-1, 1)$ , Therefore, from the formulas (39)-(41), we get the limit

$$\lim_{n \rightarrow \infty} (x_n, y_n, z_n) = (1, 1, 1), \text{ as } n \rightarrow \infty,$$

(ii) If  $x_0, y_0, z_0 \in (0, 1) \cup (1, \infty) \setminus F$ , then

$f(x_0), f(y_0), f(z_0) \in (-\infty, -1) \cup (1, \infty)$ , Therefore, from the formulas (39)-(41), we get

$$\lim_{n \rightarrow \infty} (x_n, y_n, z_n) = (-1, -1, -1).$$

(iii) If  $x_0, y_0, z_0 \in (-\infty, \infty) \setminus F$ , then we can not say about the quantities  $f(x_0)$ ,  $f(y_0)$  and  $f(z_0)$  exactly. But, the sequences

$$S_n^{(1)} = \left( \frac{1+x_0}{1-x_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+y_0}{1-y_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+z_0}{1-z_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+x_i}{1-x_i} \right)^2 \left( \frac{1-y_{i+1}}{1+y_{i+1}} \right),$$

$$S_n^{(2)} = \left( \frac{1+x_0}{1-x_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+y_0}{1-y_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+z_0}{1-z_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+y_i}{1-y_i} \right)^2 \left( \frac{1-z_{i+1}}{1+z_{i+1}} \right)$$

and

$$S_n^{(3)} = \left( \frac{1+x_0}{1-x_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+y_0}{1-y_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+z_0}{1-z_0} \right)^{\frac{2^{6n+i}}{3}} \left( \frac{1+z_i}{1-z_i} \right)^2 \left( \frac{1-x_{i+1}}{1+x_{i+1}} \right)$$

tend to either to 0 or to  $\infty$ . So, by the formulas (39)-(41), we obtain the limit

$$\lim_{n \rightarrow \infty} (|x_n|, |y_n|, |z_n|) = (1, 1, 1).$$

which completes the proof.

**References**

1. R. Abu-Saris, C. Cinar and I. Yalcinkaya, On the asymptotic stability of , Computers & Mathematics with Applications, 56(5) (2008), 1172-1175.
2. I. Dekkar, N. Touafek and Y. Yazlik, Global stability of a third-order nonlinear system of difference equations with period-two coefficients, Revista de la real academia de ciencias exactas, físicas y naturales. Serie A. Matemáticas, Doi:10.1007/s13398-016-0297-z.
3. E.M. Elsayed, Qualitative behavior of a rational recursive sequence, Indagationes Mathematicae 19(2)(2008), 189-201.
4. E.M. Elsayed, Qualitative properties for a fourth order rational difference equation, Acta Applicandae Mathematicae 110(2)(2010), 589-604.
5. N. Haddad, N. Touafek and J. F. T. Rabago, Solution form of a higher-order system of difference equations and dynamical behavior of its special case, Mathematical Methods in the Applied Sciences, (2016), doi: 10.1002/mma.4248.
6. N. Haddad, N. Touafek and J. F. T. Rabago, Well-defined solutions of a system of difference equations, Journal of Applied Mathematics and Computing, (2017), Doi:10.1007/s12190-017-1081-8.
7. Y. Halim, N. Touafek and Y. Yazlik, Dynamic behavior of a second-order nonlinear rational difference equation, Turkish Journal of Mathematics 39(6)(2015), 1004-1018.
8. A. S. Kurbanli, C. Çinar and D. Şimşek, On the periodicity of solutions of the system of rational difference equations, Applied Mathematics, 2 (2011), 410-413.
9. S. Stevic, On some solvable systems of difference equations, Applied Mathematics and Computation, 218 (2012) 5010-5018.
10. S. Stevic, M. A. Alghamdi, N. Shahzad and D. A. Maturi, On a class of solvable difference equations, Abstract and Applied Analysis, Vol. 2013, Article ID 157943, 7 pages.

11. S. Stevic, Representation of solutions of bilinear difference equations in terms of generalized Fibonacci sequences, *Electronic Journal of Qualitative Theory of Differential Equations*, 2014, No. 67, (2014), 1–15.
12. S. Stevic, J. Diblík, B. Iricanin and Z. Šmarda, On a solvable system of rational difference equations, *Journal of Difference Equations and Applications*, 20(5–6)(2014): 811–825.
13. S. Stevic, M. A. Alghamdi, A. Alotaibi and E. M. Elsayed, Solvable product–type system of difference equations of second order, *Electronic Journal of Differential Equations*, 2015, No:169, (2015), 1–20.
14. S. Stevic, New class of solvable systems of difference equations, *Applied Mathematics Letters*, 63(2017), 137–144.
15. N. Taskara, K. Uslu and D. T. Tollu, The periodicity and solutions of the rational difference equation with periodic coefficients, *Computers & Mathematics with Applications*, 62 (2011), 1807–1813.
16. D. T. Tollu, Y. Yazlik and N. Taskara, On the solutions of two special types of Riccati difference equation via Fibonacci numbers, *Advances in Difference Equations*, 2013, 2013:174.
17. D. T. Tollu, Y. Yazlik and N. Taskara, On fourteen solvable systems of difference equations, *Applied Mathematics and Computation*, 233 (2014), 310–319.
18. N. Touafek, On a second order rational difference equation, *Hacettepe Journal of Mathematics and Statistics*, 41(6) (2012), 867–874.
19. I. Yalcinkaya, C. Cinar and D. Simsek, Global asymptotic stability of a system of difference equations, *Applicable Analysis*, 87(6)(2008), 677–687, DOI: 10.1080/00036810802140657.
20. I. Yalcinkaya, On the global asymptotic stability of a second-order system of difference equations, *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 860152, 12 pages, 2008. doi:10.1155/2008/860152.
21. I. Yalcinkaya, On the global asymptotic behavior of a system of two nonlinear difference equations, *Ars Combinatoria*, 95 (2010), 151–159.
22. I. Yalcinkaya and C. Cinar, Global asymptotic stability of two nonlinear difference equations, *Fasciculi Mathematici*, 43 (2010), 171–180.
23. I. Yalcinkaya and D. T. Tollu, Global behavior of a second-order system of difference equations, *Advanced Studies in Contemporary Mathematics*, 26(4)(2016), 653–667.
24. Y. Yazlik, On the solutions and behavior of rational difference equations, *Journal of Computational Analysis & Applications*, 17(3)(2014), 584–594.
25. Y. Yazlik, E. M. Elsayed and N. Taskara, On the behaviour of the solutions of difference equation systems, *Journal of Computational Analysis & Applications*, 16(5)(2014), 932–941.
26. Y. Yazlik, D. T. Tollu and N. Taskara, On the behaviour of solutions for some systems of difference equations, *Journal of Computational Analysis & Applications*, 18(1)(2015), 166–178.
27. Y. Yazlik, D. T. Tollu and N. Taskara, On the solutions of a max–type difference equation system, *Mathematical Methods in the Applied Sciences*, 38(17)(2015), 4388–4410.
28. Y. Yazlik, D. T. Tollu and N. Taskara, On the solutions of a three–dimensional system of difference equations, *Kuwait Journal of Science* 43(1)(2016), 95–111.
29. L. Xianyi and Z. Deming, Global asymptotic stability in a rational equation, *Journal of Difference Equations and Applications*, 9(9)(2003), 833–839.
30. L. Xianyi and Z. Deming, Two rational recursive sequences, *Computers & Mathematics with Applications*, 47 (2004) 1487–1494.
31. X. Wang and Z. Li, Global asymptotic stability for two kinds of higher order recursive sequences, *Journal of Difference Equations and Applications*, 22(10)(2016), 1542–1553, DOI: 10.1080/10236198.2016.1216111.