

Mehmet ERGİN¹
Mehmet Akif KARAMERCAN²
Mehmet AYRANCI³
Yücel YAVUZ⁴
Özcan YAVAŞI⁵
Mustafa SERİNKEN⁶
Tarık ACAR⁷
Mücahit AVCİL⁸
Behçet AL⁹
Atıf BAYRAMOĞLU¹⁰
Hasan Mansur DURGUN¹¹
Yalçın GÖLCÜK¹²
İbrahim ARZIMAN¹³
Zerrin Defne DÜNDAR¹

Correspondance

Mehmet ERGİN
Necmettin Erbakan University Meram Faculty of Medicine,
Emergency Medicine Department, KONYA

Phone: 03322236778
e-mail: drmehmetergin@gmail.com

Received: 22/08/2015

Accepted: 07/10/2015

- 1 Necmettin Erbakan University Meram Faculty of Medicine, Emergency Medicine Department, KONYA
- 2 Gazi University Faculty of Medicine, Emergency Medicine Department, ANKARA
- 3 Göztepe Training and Research Hospital, Emergency Medicine Clinic, İSTANBUL
- 4 Ondokuz Mayıs University Faculty of Medicine, Emergency Medicine Department, SAMSUN
- 5 Recep Tayyip Erdoğan University Training and Research Hospital, Emergency Medicine Department, RİZE
- 6 Pamukkale University Faculty of Medicine, Emergency Medicine Department, DENİZLİ
- 7 Ordu University Training and Research Hospital, Emergency Medicine Department, ORDU
- 8 Adnan Menderes University Faculty of Medicine, Emergency Medicine Department, AYDIN
- 9 Gaziantep University Faculty of Medicine, Emergency Medicine Department, GAZİANTEP
- 10 Atatürk University Faculty of Medicine, Emergency Medicine Department, ERZURUM
- 11 Dicle University Faculty of Medicine, Emergency Medicine Department, DİYARBAKIR
- 12 Celal Bayar University Faculty of Medicine, Emergency Medicine Department, MANİSA
- 13 Gülhane Military Academy of Medicine, Emergency Medicine Department, ANKARA



RESEARCH

EPIDEMIOLOGICAL CHARACTERISTICS OF GERIATRIC PATIENTS IN EMERGENCY DEPARTMENTS: RESULTS OF A MULTICENTER STUDY

ABSTRACT

Introduction: The increasing proportion of elderly individuals in the population due to increased life expectancy has necessitated greater provision of health care. Here we aimed to determine patient characteristics, reasons for referral, and outcomes of emergency department visits and hospitalization in patients aged ≥65 years with referrals to emergency departments.

Materials and Method: This prospective, multicenter observational study was conducted over one week at the emergency departments of 13 Turkey hospitals. All patients aged ≥65 years who were referred to emergency departments with acute medical or surgical issues during the study period were included. Patients aged <65 years or those referred for trauma were excluded.

Results: In total, 1299 patients with a mean age of 74.8±7.3 years were included. Of these, 51.9% (n=674) were aged 65–74 years, 67.5% (n=877) were discharged from the hospital, and 5.8% (n=75) died during admission. The most frequently diagnosed disorders in the emergency departments were cardiovascular, gastrointestinal, and pulmonary diseases. A significant difference in age was observed between the survival and non-survival groups (p=0.001), with no significant differences in gender distribution (p=0.259), length of stay in intensive care units (p=0.605), or length of stay in hospital (p=0.055).

Conclusion: With an increased proportion of elderly individuals in the general population, the number of elderly patients referred to emergency departments continues to increase. This study presents the demographic features and clinical course of elderly patients referred to study centers.

Key Words: Elderly; Geriatric Assessment; Demography; Emergency Treatment.



ARAŞTIRMA

ACİL SERVİSLERDEKİ GERİATRİK HASTALARIN EPİDEMİYOLOJİK ÖZELLİKLERİ: ÇOK MERKEZLİ ÇALIŞMA SONUÇLARI

Öz

Giriş: Beklenen yaşam süresinin uzamasının sonucu olarak yaşlı popülasyondaki artış, bu yaş grubu için daha sık sağlık bakımı verilmesini zorunlu kılmaktadır. Bu çalışmada acil servise başvuran 65 yaş ve üzeri hastaların genel özelliklerini, başvuru nedenlerini, acil servis ve hastane ziyaretlerinin sonuçlarının saptanması amaçlandı.

Gereç ve Yöntem: Çok merkezli, prospektif, gözlemsel çalışma Türkiye’de 13 hastanenin acil servislerinde bir hafta süre ile gerçekleştirildi. Çalışma süresi içinde akut tıbbi veya cerrahi sorunlar ile acil servise başvuran 65 yaş ve üstü hastalar çalışmaya dahil edildi. Altmış beş yaş altı ve/veya travma nedeniyle başvurular ise çalışma kapsamına alınmadı.

Bulgular: Ortalama yaşı 74.8±7.3 yıl olan toplam 1299 hasta çalışmaya dahil edildi. Bu hastalardan %51.9’u (n=674) 65-74 yaş grubundaydı, %67.5’u (n=877) hastaneden taburcu edildi ve %5.8’i (n=75) yatış süreci içinde öldü. Acil serviste en sık konulan tanılar kardiyovasküler, gastrointestinal ve solunum hastalıklarıydı. Hastaneden taburcu olan ve hastane yatışı sırasında ölen hasta grupları kıyaslandığında yaş açısından istatistiksel olarak anlamlı fark varken (p=0.001), cinsiyet dağılımı (p=0.259), hastane yatış süresi (p=0.259) ve yoğun bakım ünitesi yatış süresi (p=0.055) açısından fark tespit edilmedi.

Sonuç: Yaşlı nüfusunun ve genel nüfusa oranının artışı ile birlikte yaşlı hastaların acil servis başvuru sayısı artıyor ve gelecekte daha da artacaktır. Bu çalışma, çalışma merkezlerine başvuran yaşlı hastaların demografik özelliklerini ve klinik seyirlerinin sonuçlarını ortaya koymaktadır.

Anahtar Sözcükler: Yaşlı; Geriatrik Değerlendirme; Demografi; Acil Tedavi.



INTRODUCTION

According to 2014 population statistics, citizens aged ≥ 65 years comprise 8% of the total population in Turkey. This ageing population comprises over 6 million people, surpassing the total population of a number of countries, including Denmark, Slovakia, Finland, and Norway. Furthermore, this age group comprises 8.3% of the total global population. Among 228 countries, Monaco ranks first with the largest elderly population, followed by Japan and Germany, whereas Turkey ranks 94th. It has been predicted that the proportion of elderly individuals in Turkey will be 10.2% of the total population in 2023, 20.8% in 2050, and 27.7% in 2075 (1). The increased proportion of elderly individuals is predominantly due to increased life expectancy, resulting in a range of associated health issues that necessitate greater provision of health care.

When compared with young individuals, elderly patients are referred to emergency departments (EDs) more often and with more complicated presentations, require longer duration in intensive care units (ICUs) and EDs, and are more frequently hospitalized (2-6). Hence, health professionals working in EDs should be familiar with the characteristics of elderly patients and common reasons for referral in order to provide the necessary infrastructure and training. Thus, a greater level of expertise in the care for the elderly can be attained. The purpose of this multicenter study was to determine patient characteristics, reasons for referral, and outcomes of ED visits and hospitalizations in patients aged ≥ 65 years with ED referrals.

MATERIALS AND METHOD

Study Design

This prospective, multicenter observational study was conducted over one week at the EDs of 13 hospitals belonging to universities, state, and military. The study included patients aged ≥ 65 years with medical or surgical issues requiring admission to EDs. The purpose of the present study was to determine patient characteristics and the outcomes of ED visits in a geriatric population in Turkey.

Study Settings

This study was conducted in 13 hospitals in the provinces of Ankara, Aydin, Denizli, Diyarbakir, Erzurum, Gaziantep, Istanbul, Konya, Manisa, Ordu, Rize, and Samsun in 7 different geographical regions of Turkey between September 1 and 7,

2014. All hospitals were academic centers with emergency medicine specialist training programs. The coordination centers of the present study were Konya and Ankara. All patients aged ≥ 65 years who were referred to EDs with acute medical or surgical issues during the study period were included in the present study. Patients aged < 65 years or those referred for trauma were excluded. Patients who had received pre-hospital cardiopulmonary resuscitation or were transferred to hospitals other than the study centers were also excluded from the present study.

Study Protocol

The following data for patients who met the study inclusion criteria were recorded: age, gender, major complaints, comorbidities, medications, diagnosis, admission and discharge information, admission wards, length of stay (LOS) in ICUs, LOS in hospital, and in-hospital mortality. Patients were followed-up until discharge from the hospital or death.

Statistical Analyses

Statistical Package for the Social Science Version 16.0 (SPSS Inc., Chicago, IL, USA) was used for the statistical analysis of all obtained data. Categorical data were expressed as numbers and percentage and continuous data as mean \pm standard deviation (SD). Univariate analyses were performed using the Chi-square test for categorical variables and Student's *t*-test for continuous variables. P-values of < 0.05 were considered statistically significant.

RESULTS

A total of 1299 patients aged ≥ 65 years referred to EDs with non-traumatic, acute surgical, or medical problems were included in the present study. The mean age of the included patients was 74.8 ± 7.3 years. Of these, 674 (51.9%) were aged 65–74 years and 145 (11.2%) were aged ≥ 85 years, with 680 (52.3%) females (Table 1). No significant differences in gender distribution were observed between the age groups ($p=0.422$) (Table 2).

The most frequent complaints were related to gastrointestinal (25.8%, $n=333$), pulmonary (18.2%, $n=237$), and central nervous (16.8%, $n=218$) systems (Table 1). No significant difference in the distribution of complaints was observed between the age groups ($p=0.081$).

According to medical histories provided by patients and/or their relatives, 10.2% ($n=133$) of patients had no previous history of a known disease. The most frequent comorbi-



Table 1— Distribution of Demographic Features, Leading Complaints, Comorbidities and Number of Daily Medications, Results of ED and Hospital Visiting of the Study Group

A. Demographic parameters		D. Number of Medications^β	
Age ^α	74.8±7.3	2	282 (21.7%)
Age groups ^β		4-6	254 (19.6%)
65-74	674 (51.9%)	Any known medication	251 (19.3%)
75-84	480 (37.0%)	1	251 (19.3%)
≥85	145 (11.2%)	3	227 (17.5%)
Total	1299 (100%)	>6	34 (2.6%)
Gender ^β		Total	1299 (100%)
Female	680 (52.3%)	E. ED and hospital visiting results	
Male	619 (47.7%)	ED visiting ^β	
Total	1299(100.0%)	Discharged from ED	877 (67.5%)
B. Leading Complaints^{βλ}		Admission to a ward	282 (21.7%)
Gastrointestinal	330 (25.4%)	Admission to the ICU	140 (10.8%)
Pulmonary	237 (18.2%)	Total	1299 (100.0%)
Central nervous system	218 (16.8%)	Length of stay in hospital ^α	8.3 ± 6.7
Multisystem	194 (14.9%)	Length of stay in ICU ^α	6.3 ± 6.9
Cardiovascular	125 (9.6%)	Exitus during hospital stay ^β	75 (5.8%)
Skeletomuscular	96 (7.4%)	F. Last Diagnosis^{βλ}	
Genitourinary	60 (4.6%)	Cardiovascular	253 (19.5%)
Others	39 (3.0%)	Gastrointestinal	228 (17.6%)
Total	1299(100.0%)	Pulmonary	197 (15.2%)
C. Comorbidities^{βΩ}		Genitourinary	98 (7.5%)
Hypertension	673 (51.8%)	Neurology	95 (7.3%)
Diabetes mellitus	321 (24.7%)	Others	74 (5.7%)
Coronary heart disease	279 (21.5%)	Infection	73 (5.6%)
COPD / asthma	232 (17.9%)	Nose-throat-ear	68 (5.2%)
Malignity	198 (15.2%)	Skeletomuscular	65 (5.0%)
Congestive heart failure	150 (11.5%)	Oncology	54 (4.2%)
Any known disease	133 (10.2%)	Nephrology	45 (3.5%)
Cerebrovascular accident	84 (6.5%)	Endocrine	22 (1.7%)
Chronic renal failure	79 (6.1%)	Psychiatric	17 (1.3%)
Hyperlipidemia	78 (6.0%)	Intoxication	10 (0.8%)
Arrhythmia	70 (5.4%)	Total	1299 (100.0%)
Dementia / Alzheimer disease	64 (4.9%)		

ED: Emergency Department; ICU: Intensive Care Unit; ^αDescribed as mean value ± SD and year; ^βDescribed as number (n) and percentage (%);

^αDescribed as mean value ± SD and day; ^λClassified according to the organ system; ^ΩOne patient can have more than one disease

dities in the remaining patients were hypertension (51.8%, n=673), diabetes mellitus (24.7%, n=321), and coronary heart disease (21.5%, n=279). Of these patients, 19.3% (n=251) had no history of regular medication usage and 61.4% (n=797) had a history of multiple medication usage (two or more medications per day) (Table 1).

ED referrals ended in discharge for 67.5% (n=877) of patients and hospitalization in a ward or ICU in 32.1% (n=422)

of patients (Table 1). A significant difference in the rate of discharge or admission was observed between the age groups (p=0.034). The rate of ED discharge was higher in the 65–74 age group (71.2%) and the rates of admission to a ward or ICU were higher in the 75–84 age group (25.2% and 12.5%, respectively) (Table 2).

The most frequently diagnosed disorders in patients referred to EDs were cardiovascular (19.5%, n=253), gastrointes-



Table 2— Distribution of Parameters by Age Groups

Parameters	Age Group	65 - 74 (n=674)	75 - 84 (n=480)	≥85 (n=145)	p value
Gender ^β	Male	322 (47.8%)	235 (49%)	62 (42.8%)	0.422
	Female	352 (52.2%)	245 (51%)	83 (57.2%)	
Result of ED visiting ^β	Patients discharged from ED	480 (71.2%)	299 (62.3%)	98 (67.6%)	0.034
	Patients admitted to a ward	131 (19.4%)	121 (25.2%)	30 (20.7%)	
Patients admitted to the ICU		63 (9.3%)	60 (12.5%)	17 (11.7%)	
Length of stay in ICU ^α		6.73±7.56	6.60±6.90	5.21±6.35	0.683
Length of stay in hospital ^α		10.33±8.80	8.06±7.82	7.53±6.29	0.340
Result of hospital visiting ^β	Patients discharged from hospital	646 (95.8%)	448 (93.3%)	130 (89.7%)	0.009
	In hospital mortality	28 (4.2%)	32 (6.7%)	15 (10.3%)	

ED: Emergency Department; ICU: Intensive Care Unit; ^βDescribed as number (n) and percentage (%); ^αDescribed as mean value ± SD and day.

tinal (17.6%, n=228), and pulmonary (15.2%, n=197) diseases (Table 1). No significant differences in the distribution of final diagnoses was observed between the age groups (p=0.067).

Patients were most frequently hospitalized in internal medicine (28.5%, n=119), cardiology (19.4%, n=81), and respiratory disease (13.4%, n=56) wards. The mean LOS in hospital was 8.3 ± 6.7 days, whereas the mean LOS in ICUs was 6.3±6.9 days. Of the admitted patients, 75 (5.8%) died during hospitalization (Table 1). No statistically significant difference in the LOS in ICUs or hospital were observed between

the age groups (p=0.683, and p=0.340, respectively). However, a significant difference in the prognosis of hospitalized patients was observed between the age groups (p=0.009). The 65–74 age group had a higher rate of discharge (95.8%) compared with the ≥85 age group, which had a high mortality rate (10.3%) (Table 2).

Patients referred to EDs were categorized into “survival” and “non-survival” groups (Table 3). A significant difference in age (p=0.001) was observed between the survival and non-survival groups, with no significant difference in gender distribution (p=0.259), LOS in ICUs (p=0.605), and LOS in

Table 3— Distribution of Parameters by Survival and Non-survival Groups

Parameters	Groups	Survival (n=1224)	Non-survival (n=75)	p value
Age ^α		74.64±7.26	77.45±7.47	0.001
Gender ^β	Male	588 (48%)	31 (41.3%)	0.259
	Female	636 (52%)	44 (58.7%)	
Length of stay in ICU ^α		6.07±6.9	6.73±6.85	0.605
Length of stay in hospital ^α		8.55±6.68	6.75±6.59	0.055
Results of hospital admissions (n=422) ^β	Patients admitted to a ward	256 (73.8%)	26 (34.7%)	<0.001
	Patients admitted to the ICU	91 (26.2%)	49 (65.3%)	

ED: Emergency Department; ICU: Intensive Care Unit; ^αDescribed as mean value ± SD and year; ^βDescribed as number (n) and percentage (%); ^αDescribed as mean value ± SD and day.



hospital ($p=0.055$). When the 422 patients who were hospitalized following referral to ED were evaluated, a significant difference in mortality was observed between patients hospitalized in wards and patients admitted to ICUs ($p<0.001$). A higher mortality rate was observed in patients admitted to ICUs (65.3%) (Table 3). A significant difference in mortality was also observed according to the major complaint requiring ED referral ($p=0.013$). The non-survival group had higher rates of multisystemic (29.3%, $n=22$), pulmonary (21.3%, $n=16$), and gastrointestinal (20%, $n=15$) complaints. Furthermore, a significant difference in mortality was observed according to the final diagnosis made in the ED ($P<0.001$). In the non-survival group, higher rates of diagnosed pulmonary (21.3%, $n=16$), nephrologic (21.3%, $n=16$), and infectious (17.3%, $n=13$) diseases were observed.

DISCUSSION

In Turkey, 3.6%–41% of patients referred to EDs are elderly (2-4, 7-9). In developed countries, 20% of referrals to EDs are patients aged ≥ 65 years (10-12). These rates are a reflection of the ageing populations of developed countries.

Previous single-center studies in Turkey have reported the mean age of elderly referrals to EDs as 74.50 ± 6.53 (4), 73.18 ± 6.94 (7), and 74.5 ± 8.5 (13) years. The mean age of patients in the present study is consistent with these previous findings in Turkey (4,7,13). The mean age of elderly patients referred to EDs has been reported as 76.8 ± 8.0 years by an Irish study (10) and 82 years (min, 60; max, 103) by a Chinese study (14).

In the present study, we observed a greater number of ED referrals in the 65–74 age group compared with other age groups. A previous study reported more referrals in the 65–74 age group, whereas another study reported the highest number of ED referrals in the 75–84 age group (3,4). The findings of the present study are consistent with those of a previous study reporting that 60.3% of the aged population in Turkey is in the 65–74 age group (1).

The female/male ratio among elderly groups referred to EDs is reportedly variable, with a female predominance. In Turkey, the proportion of female patients among elderly patients with ED referrals reportedly ranges from 44.2%–59.2% (3,4,7,13). In the present study, more females than males were observed in all elderly patient groups. Consistent with the results of Kapçi *et al.* (3), no difference in gender distribution was observed between the age groups. These findings may be due to the greater number of females than

males in the geriatric population in Turkey (1). A Chinese study (14) reported that more males are referred to EDs, whereas an Irish study (10) reported that more females are referred.

In Turkey, the prevalence of chronic diseases among elderly patients ranges from 72.6%–94.4% (15,16). It has been reported that among elderly patients referred to EDs, 72.8% have at least 1 and 32.5% have ≥ 2 chronic disease(s) (13). Consistent with the results of the present study, Loğoğlu *et al.* (4) reported that the most frequently observed disorders in elderly patients referred to EDs are hypertension, coronary artery disease, and diabetes mellitus.

The increased incidence of a number of disorders with advanced age and the tendency to prescribe multiple medications to patients staying in residential services have changed the pattern of medication use in elderly patients (17). Özşaker *et al.* (13) reported that 56.2% of elderly patients referred to EDs were receiving continuous pharmacological treatment. We observed a high prevalence of chronic disease and frequent continuous pharmacological treatment in elderly patients referred to EDs. In light of these findings, health professionals in EDs should be aware of the different pharmacokinetic and pharmacodynamic features of particular medications and consequent variations in pharmacological response among elderly patients (18). Furthermore, doctors working in EDs should determine the medications currently being used by elderly patients and consider these during diagnosis and treatment planning.

The most frequent complaints among elderly patients referred to EDs in Turkey are related to the respiratory, gastrointestinal, cardiovascular, and central nervous systems (3,4, 7,8,13), findings corroborated by the present study.

It has been reported that the most frequent diagnoses following referral to EDs are pneumonia, cerebrovascular disease, and chronic obstructive pulmonary disease (2). On the other hand, Kapçi *et al.* (3) reported that gastrointestinal, respiratory, and neurological diseases are the predominant disorders in such patients. In the present study, diagnoses were most frequently related to cardiovascular, gastrointestinal, and respiratory systems (Table 1). The most frequently reported disorders among elderly patients with ED referrals are respiratory, cardiological, and neurological diseases in China (14); infectious diseases, predominantly malaria, and hypertension-related cardiovascular diseases in Nigeria (19); and infectious, cardiovascular, and gastrointestinal diseases in Ireland (10).



The hospitalization rate of elderly patients referred to EDs in Turkey ranges from 11.5%–47.2%. Of the elderly patients hospitalized, 65.4%–81.4% are hospitalized in wards and 18.6%–34.6% in ICUs. These patients are most frequently hospitalized in cardiology, internal medicine, respiratory disease, neurology, and general surgery wards (3,4,7,13). An Irish study (10) reported the rate of hospitalization among elderly patients referred to ED as 63.5%. In the present study, we observed a higher rate of hospitalization, with patients most frequently hospitalized in internal medicine, cardiology, and respiratory disease wards. The results of the present study indicate elderly patients in Turkey are predominantly referred to EDs due to medical complaints rather than surgical complaints, with the exception of trauma.

LOS in hospital among elderly patients varies among previous studies. Regarding the mean LOS in hospital, Baz *et al.* (7) have reported 7.1±6.2 days whereas Loğoğlu *et al.* (4) have reported 4±6.6 days. In the present study, we observed a mean LOS in hospital of 8.3±6.7 days. Kennely *et al.* (10) reported a mean LOS in hospital of 13.1±19.0 days and found that the only important multivariate predictor of prolonged hospitalization was advanced age. We observed no differences in LOS in hospital or ICUs between the age groups in the present study (Table 1).

Among patients included in the present study, 5.8% (n=75) died during hospital admission. Single-center studies of elderly patients in Turkey have reported that following referral to EDs, the mortality rate in the ED ranges between 0.1% and 7.6% and in hospital ranges between 0.8% and 11.5% (7,8). A previous Irish study (10) reported a mortality rate in the ED of 14.6% in elderly patients. In contrast with the results of the present study, Üstündağ *et al.* (8) found no difference in age distribution between patients who survived compared with those who did not. The same authors (8), in a finding consistent with our results, found no difference in LOS in hospital. Further, this study reported that altered level of consciousness, respiratory distress, and abdominal pain are more frequent among patients referred to EDs who do not survive (8). In contrast with our results, according to data from the Turkish Statistics Institute (1), the most frequent causes of mortality in elderly patients are cardiovascular system disorders (46.8%), benign and malignant tumors (17.7%), and respiratory system disorders (11.7%). Many scoring systems for the prediction of hospitalization and prognosis have been proposed. Clinical studies have reported that the Modified Early Warning Score (MEWS) and VitalPAC Early Warning Score (VIEWS) have efficacy in the prediction of

hospitalization and mortality during hospital stays and that the Rapid Emergency Medicine Score (REMS), REMS without the inclusion of age, and HOTEL scoring systems (Hypotension, Oxygen saturation, low Temperature, ECG changes, Loss of independence) have efficacy in the prediction of ICU stay duration and mortality during hospital stays (20,21). It has been posited that high scores on the Charlson Comorbidity Index indicate a high risk of mortality during hospital admissions (10). Mortality rates are dependent on patient factors as well as the quality of health care. Factors associated with mortality should be assessed and those that can be modified should be corrected. It has been reported that the use of the “Comprehensive Geriatric Assessment (CGA)” increases survival by up to 12 months in elderly patients referred to hospitals. The Comprehensive Geriatric Assessment can also be used in EDs (22,23).

We believe that the discrepancy between the results of the present study and those of previous studies are related to the frequency and reasons for the referral of elderly patients to EDs. Furthermore, subsequent clinical courses and prognoses have been shown to be related to the degree of development and demographic, socio-cultural, and climate differences among countries as well as health care infrastructure, diagnostic capability in treatment centers, and different approaches to the treatment of patients.

Concomitant with an increased proportion of elderly individuals in the general population, the number of elderly patients referred to EDs continues to increase. This multicenter study presents the demographic features and clinical course of elderly patients referred to EDs. The findings of the present study may guide the planning of future studies on the measurement and improvement of health care provided to elderly patients by EDs in Turkey.

Conflict of Interest

All authors have no conflict of interest to declare.

REFERENCES

1. Statistics Institute of Turkey. Elderly statistics 2014. In: Birol Aydemir (Ed). 1st edition, TÜİK Press, Ankara, Türkiye.2015, pp:1-91. [Internet] Available from:http://www.tuik.gov.tr/Kitap.do?Fmetod=KitapDetay&KT_ID=11&KITAP_ID=265. Accessed:01.06.2015 (in Turkish).
2. Günalp M, Gülünay B, Polat O, et al. Increased length of stay in emergency department in Turkey: due to inappropriate emergency department use or aging? *Turkiye Klinikleri J Med Sci* 2014;34(3):273-9. (DOI: 10.5336/medsci.2013-34423).



3. Kapıcı M, Tomruk Ö, Beceren NG, et al. Investigation of factors affecting cost of geriatric patients admitted to the emergency department. *JAEM* 2013;12(3):134-8.
4. Loğoğlu A, Ayırık C, Köse A, et al. Analysis of nontraumatic elderly patient presentations to the emergency department. *Tr J Emerg Med* 2013;13(4):171-79. (in Turkish).
5. Karadağ B, Çat H, Öztürk AO, Basat O, Altuntaş Y. Patients admitted to the emergency outpatient clinic and kept under observation. *Journal of Academic Geriatrics* 2010;2(3):176-85. [Internet]? Available from: http://www.akadgeriatri.org/managete/fu_folder/2010-03/html/2010-2-3-176-185.htm Accessed:01.06.2015. (in Turkish).
6. Taymaz T. Examination of geriatric patients hospitalized from the emergency department. *Journal of Academic Geriatrics* 2010;2(3):167-75. [Internet] Available from: http://www.akadgeriatri.org/managete/fu_folder/2010-03/html/2010-2-3-167-175.htm Accessed:01.06.2015. (in Turkish).
7. Baz Ü, Satar S, Kozacı N, Açıklın A, Gülen M, Karakurt Ü. Geriatric patient admissions to the emergency service. *JAEM* 2014;13:53-7.
8. Üstündağ M, Tunç İ, Orak M, Özhasenekler A, Durgun HM, Güloğlu C. Clinical and demographic characteristics of patients aged 65 years and older admitted to emergency department. *Journal of Academic Geriatrics* 2013;5(3):117-26. [Internet] Available from: http://www.akadgeriatri.org/managete/fu_folder/2013-03/2013-5-3-117-126.pdf Accessed:01.06.2015.
9. Şahin S, Boydak B, Savaş S, Yalçın A, Akçiçek F. Characteristics of patients aged 65 and over in the emergency department. *Journal of Academic Geriatrics* 2011;3(1):41-6. [Internet] Available from: http://www.akadgeriatri.org/managete/fu_folder/2011-01/2011-3-1-041-046.pdf Accessed:01.06.2015. (in Turkish).
10. Kennelly SP, Drumm B, Coughlan T, Collins R, O'Neill D, Romero-ortuno R. Characteristics and outcomes of older persons attending the emergency department: a retrospective cohort study. *QJM* 2014;107(12):977-87. (PMID:24935811).
11. Xu KT, Nelson BK, Berk S. The changing profile of patients who used emergency department services in the United States. 1996 to 2005. *Ann Emerg Med* 2009;54(6):805-810.e1-7 (PMID:19811852).
12. Roberts DC, McKay MP, Schaffer A. Increasing rates of emergency department visits for elderly patients in the United States, 1993 to 2003. *Ann Emerg Med* 2008;51(6):769-74. (PMID:18069088).
13. Özşaker E, Korkmaz FD, Dölek M. Analyzing individual characteristics and admission causes of elderly patients to emergency departments. *Turkish Journal of Geriatrics* 2011;14(2):128-34. [Internet] Available from: http://geriatri.dergisi.org/pdf/pdf_TJG_570.pdf Accessed:01.06.2015. (in Turkish).
14. Liu HW, Han LN, Zhao YX, Zhang W. Common causes of geriatric medical emergencies in China. *J Geriatr Cardiol* 2015;12(1):91-92. (PMID:25678910).
15. Ünsaldı ÜE, Piyal B. Evaluating the chronic diseases and activity restriction in a group of subjects of aged 65 years and over that applied to Çubuk Health Center. *Turkiye Klinikleri J Med Sci* 2002;22(4):362-8. [Internet]? Available from: <http://www.turkiyeklinikleri.com/article/tr-cubuk-saglik-ocagina-basvuran-65-yas-ve-uzeri-grupta-kronik-hastalik-ve-etkinlik-kisitiliginin-degerlendirilmesi-2464.html> Accessed:01.06.2015. (in Turkish).
16. Özçakar N, Kartal M, Güldal D. Medical students visits to a nursing home: what does it add to quality of life? *Turkiye Klinikleri J Med Sci* 2010;30(1):17-23. (in Turkish).
17. Giron MST, Claesson C, Thorslund M, Oke T, Winbland B, Fastbom J. Drug use patterns in a very elderly population: a seven year review. *Clin Drug Invest* 1999;17(5):398-98. [Internet] Available from: <http://link.springer.com/article/10.2165/00044011-199917050-00005> Accessed:01.06.2015.
18. Göksel SÜ. Drug use in elderly and integrative medicine. *Turkiye Klinikleri J Med Sci* 2009;29(suppl):S76-S79. [Internet] Available from: <http://www.turkiyeklinikleri.com/article/tr-yaslida-ilac-kullanimi-ve-butunleyici-tip-56071.html> Accessed:01.06.2015. (in Turkish).
19. Iloh G, Amadi A, Awa-Madu J. Common geriatric emergencies in a rural hospital in South - Eastern Nigeris. *Niger J Clin Pract* 2012;15(3):333-7. (PMID:22960971).
20. Dundar ZD, Ergin M, Karamercan MA, et al. Modified early warning score and VitalPac Early warning score in geriatric patients admitted to emergency department. *Eur J Emerg Med* 2015 Apr 21. (PMID:25919485).
21. Dundar ZD, Karamercan MA, Ergin M, et al. Rapid emergency medicine score and HOTEL score in geriatric patients admitted to the emergency department. *International Journal of Gerontology* 2015;9(2):87-92.
22. Ellis G, Whitehead MA, O'Neill D, Langhorne P, Robinson D. Comprehensive geriatric assesment for older adults admitted to hospital. *Cochrane Database Syst Rev* 2011;6(7):CD006211. (PMID:21735403).
23. Conroy SP, Ansari K, Williams M, et al. A controlled evaluation of comprehensive geriatric assesment in the emergency department: the 'Emergency Frailty Unit'. *Age Aging* 2014;43(1):109-14. (PMID:23880143).